

DESIGNED
with you in mind



Not actual patients. Results may vary.

5 THINGS to know about your
5-DAY TREATMENT
for actinic keratosis (AK)

For more information on KLISYRI[®], please click to see [Patient Information](#) and full [Prescribing Information](#).

1

WHY YOUR HEALTHCARE PROFESSIONAL PRESCRIBED KLISYRI®

- KLISYRI® (tirbanibulin) is the **shortest topical prescription AK treatment** available today—**only 5 days**
- KLISYRI® was studied in the largest AK clinical trial program of its kind*



- **5x more** KLISYRI® patients achieved complete clearance than did placebo patients†
 - The most common side effects with KLISYRI® were local skin reactions, including itching and pain in the treatment area
 - Fewer than 1 in 10 patients had severe local skin reactions (LSRs)
 - LSRs are common, mostly mild to moderate, but can be severe during treatment with KLISYRI®
- No allergic reactions to UV rays occurred in healthy volunteers using KLISYRI® in clinical trials‡

See **“What are the possible side effects of KLISYRI®?”** in part 5 of this brochure.

*The largest pivotal clinical trial program of an AK topical prescription treatment.

†Based on a pooled analysis of data from two Phase 3 studies with KLISYRI®. 100% AK clearance rates for the two studies were 44% KLISYRI® vs 5% vehicle in study 1 and 54% KLISYRI® vs 13% vehicle in study 2.

‡Clinical studies in healthy subjects demonstrated KLISYRI® did not cause contact sensitization, phototoxic skin reactions, or photoallergic skin reactions.

AK: actinic keratosis; UV: ultraviolet.

2

WHAT IS KLISYRI®

- KLISYRI® is a Food and Drug Administration (FDA)-approved prescription medicine used on the skin to treat actinic keratosis (AK) on the face or scalp
- KLISYRI® has a unique way of working and is different from current topical AK therapies you may have already tried
- KLISYRI® is an ointment that comes in single packets you use daily for **5 days** in a row

FILLING YOUR KLISYRI® PRESCRIPTION*



If you receive a **SAMPLE**, use it only if you intend to fill your prescription the very next day

Get your **5-day** prescription pack from your pharmacy

Use a total of **5 packets**— one packet per day

Use 5 packets, 5 days in a row ONLY

***NOTE:** If you received a starter sample from your healthcare professional and then get your prescription filled at your pharmacy, you will have an extra packet of medication. Do not use a 6th packet.

- It is not known if KLISYRI® is safe and effective for children less than 18 years of age

For more information on KLISYRI®, please click to see [Patient Information](#) and full [Prescribing Information](#).

3

HOW TO FILL YOUR PRESCRIPTION

1.



Your doctor sends your **KLISYRI® (tirbanibulin)** prescription electronically to the specialty pharmacy. **There's no paper prescription.**

2.



The specialty pharmacy calls you—likely from **a toll-free number**. Be sure to pick up this call. **Your KLISYRI® prescription cannot be filled until they speak with you.**

3.



Pay no more than \$95 if you have commercial insurance coverage. Not sure if you qualify for savings? Ask the specialty pharmacy when they call.

4.



Your KLISYRI® prescription is conveniently shipped to your door.

For more information on KLISYRI®, please click to see [Patient Information](#) and full [Prescribing Information](#).

4

HOW TO APPLY KLISYRI®



1.



Open one KLISYRI® (tirbanibulin) ointment packet

2.



Place ointment on finger

3.



Apply ointment evenly.
You can cover an area of up to about 4 square inches

4.



Wash hands with soap and water immediately after application and discard the packet after use, even if you have not used entire packet

- KLISYRI® should not be used if your skin is not healed from any previous medicine, procedure, or surgical treatment
- Do not use more KLISYRI® than you need to cover the treatment area
- Using too much KLISYRI®, or using it too often or for too long can increase your chances of having a severe skin reaction or other side effects

5

OTHER IMPORTANT INFORMATION TO KNOW ABOUT USING KLISYRI®

Important: KLISYRI is for use on the skin only (topical). Do not use KLISYRI in, around, or near your eyes, mouth or lips.

What is KLISYRI?

KLISYRI is a prescription medicine used on the skin to treat actinic keratosis on the face or scalp.

It is not known if KLISYRI is safe and effective in children less than 18 years of age.

Before using KLISYRI, tell your healthcare provider about all of your medical conditions, including if you:

- are being treated or have been treated for actinic keratosis with any other medicine, procedure, or surgery. You should not use KLISYRI until your skin has healed from other treatments.
- have other skin problems in the treatment area.
- are pregnant or plan to become pregnant. It is not known if KLISYRI can harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if KLISYRI passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with KLISYRI.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements.

How should I use KLISYRI?

- Use KLISYRI as your healthcare provider tells you to. KLISYRI is for skin use only.
- Apply KLISYRI to evenly cover the treatment area(s) on the face or scalp 1 time a day for 5 days in a row (consecutive) using 1 single dose packet with each application. Do not apply KLISYRI to other areas.
- Do not use more KLISYRI than you need to cover the treatment area. Using too much KLISYRI, or using it too often, or for too long can increase your chances of having a severe skin reaction or other side effects.
- Do not cover the treatment area with a dressing after application of KLISYRI. Irritation of the skin may happen if a dressing is applied to the treatment area.

- Do not get KLISYRI in, around, or near your eyes. Do not touch your eyes while you are applying KLISYRI.
 - Wash your hands right away with water and soap after applying KLISYRI. After applying KLISYRI, be careful to keep KLISYRI on the treated area from coming into contact with your eyes. Irritation may happen if you get KLISYRI in your eyes.
 - If you accidentally get KLISYRI in your eyes, flush them with water and get medical care as soon as possible. See, “What are the possible side effects of KLISYRI?”
- Do not get KLISYRI in, around, or near your mouth or lips.
- Avoid washing and touching the treated area for approximately 8 hours after application of KLISYRI. After 8 hours, you may wash the area with a mild soap and water.
- Avoid transferring the product to other areas after application.
- Throw away any open packet of KLISYRI after use even if there is medicine still left in it.

What are the possible side effects of KLISYRI?

KLISYRI may cause serious side effects, including:

- Eye irritation can happen if KLISYRI gets into your eyes. If you accidentally get KLISYRI into your eyes, flush them with water and get medical care as soon as possible.
- Local skin reactions are common but can also be severe during treatment with KLISYRI. Call your healthcare provider if you develop any local skin reactions including redness, flaking or scaling, crusting, or swelling that is more severe, or if you get blisters, peeling, pus, ulcers, or breakdown of your skin.

The most common side effects of KLISYRI include: itching or pain in the treatment area.

These are not all of the possible side effects of KLISYRI. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

For more information on KLISYRI®, please click to see [Patient Information](#) and full [Prescribing Information](#).

DESIGNED
with you in mind

Not actual patients. Results may vary.

Visit www.klisyri.com to learn more.

For more information on KLISYRI[®], please click to see [Patient Information](#) and full [Prescribing Information](#).